

ON A CERTAIN FORMULATION OF THE EINSTEIN EQUATIONS *

PAWEŁ NUROWSKI

Dipartimento di Scienze Matematiche
Università degli Studi di Trieste, Trieste, Italy [†]

February 7, 2008

Abstract

We define a certain differential system on an open set of $\mathbf{R}^6$. The system locally defines a Lorentzian 4-manifold satisfying the Einstein equations. The converse statement is indicated and its details are postponed to the forthcoming paper.

*Research supported in part by: Komitet Badań Naukowych (Grant nr 2 P03B 017 12), Consorzio per lo Sviluppo Internazionale dell'Università degli Studi di Trieste and Erwin Schrödinger International Institute for Mathematical Physics.

[†]Permanent address: *Katedra Metod Matematycznych Fizyki, Wydział Fizyki, Uniwersytet Warszawski, ul. Hoża 74, Warszawa, Poland, e-mail: nurowski@fuw.edu.pl*

Let $\mathcal{U}$ be an open subset of $\mathbf{R}^6$. Suppose that on $\mathcal{U}$ we have six 1-forms $(\Lambda, F, \bar{F}, T, E, \bar{E})$ which satisfy the following conditions

- i) Λ, T are real- and F, E are complex-valued 1-forms
- ii) $\Lambda \wedge F \wedge \bar{F} \wedge T \wedge E \wedge \bar{E} \neq 0$ at each point p of $\mathcal{U}$
- iii) there exist complex-valued 1-forms Ω and Γ on $\mathcal{U}$, and a certain complex function α on $\mathcal{U}$ such that

$$\begin{aligned}
dF &= (\Omega - \bar{\Omega}) \wedge F + E \wedge T + \Gamma \wedge \Lambda \\
dT &= \bar{\Gamma} \wedge F + \Gamma \wedge \bar{F} - (\Omega + \bar{\Omega}) \wedge T \\
d\Lambda &= \bar{E} \wedge F + E \wedge \bar{F} + (\Omega + \bar{\Omega}) \wedge \Lambda \\
dE &= 2\Omega \wedge E + \bar{F} \wedge T + \alpha \Lambda \wedge F.
\end{aligned} \tag{1}$$

The aim of this paper is to prove that we can associate to such $\mathcal{U}$ a Lorentzian 4-manifold that satisfies the Einstein equations. The proof will be a direct consequence of the following sequence of (very simple) Lemmas.

Lemma 1

If $(\Lambda, F, \bar{F}, T, E, \bar{E})$ satisfy (1) then there exist complex functions a, h on $\mathcal{U}$ and a real constant λ such that

$$\begin{aligned}
d\Gamma &= 2\Gamma \wedge \bar{\Omega} + \bar{\alpha}T \wedge F + \bar{a}(T \wedge \Lambda - F \wedge \bar{F}) + h\Lambda \wedge \bar{F} \\
d\Omega &= E \wedge \bar{\Gamma} - (\alpha + \frac{\lambda}{2})(T \wedge \Lambda + F \wedge \bar{F}) + a\Lambda \wedge F.
\end{aligned} \tag{2}$$

The proof of this Lemma is straightforward but lengthy. It is performed by imposing conditions

$$d^2\Lambda = d^2T = d^2F = d^2E = 0 \tag{3}$$

on the system (1) and by using the independence condition **ii**). Application of (3) implies the possible forms of $d\Gamma$ and $d\Omega$. Use of

$$d^2\Gamma = d^2\Omega = 0 \tag{4}$$

shows, in particular, that λ must be a real constant. $\square$

There are also other equations that are implied by (3)-(4). They carry information

about the differentials of α, a, h and about the decompositions of Γ and Ω onto the basis 1-forms $(\Lambda, F, \bar{F}, T, E, \bar{E})$. Explicitely we have

$$\begin{aligned} d\alpha &= -2aE - \bar{\gamma}_3 T + \bar{\gamma}_1 \bar{F} + sF - w\Lambda, \\ da &= \bar{h}E - (3\alpha + \lambda)\bar{\Gamma} - 2a\Omega + sT - w\bar{F} + rF - t\Lambda, \\ dh &= -4h\bar{\Omega} + 4\bar{a}\Gamma - \bar{r}T + \bar{t}F - u\Lambda + v\bar{F}, \end{aligned} \tag{5}$$

where the possible forms of Ω and Γ are

$$\begin{aligned} \Omega &= -\omega_1\Lambda + \omega_2 F + \omega_3 \bar{F} + \omega_4 T, \\ \Gamma &= \gamma_1\Lambda + 4\bar{\omega}_1 F - \gamma_3 \bar{F} - 4\bar{\omega}_2 T - (3\bar{\alpha} + \lambda)\bar{E}. \end{aligned} \tag{6}$$

Here $\gamma_1, \gamma_3, s, w, r, t, u, v, \omega_1, \omega_2, \omega_3, \omega_4$ are certain complex functions on $\mathcal{U}$.

Note that the differential system (1), (2), (5), (6) is still not closed. For example, the equations of the sort $d^2\alpha = 0$ should be still imposed.

Lemma 2

$\mathcal{U}$ is locally foliated by 2-dimensional manifolds $\mathcal{S}_x$, which are tangent to the real distribution $\mathcal{V}$ defined by

$$\Lambda(\mathcal{V}) = F(\mathcal{V}) = T(\mathcal{V}) = 0.$$

Here the proof is an immediate application of the Froebenius theorem, since the forms $(\Lambda, F, \bar{F}, T)$ form a closed differential ideal due to the relations (1). $\square$

Let us define

$$G = F\bar{F} - T\Lambda \tag{7}$$

on $\mathcal{U}$. G constitutes a degenerate metric on $\mathcal{U}$. It has the signature $(+, +, +, -, 0, 0)$.

Lemma 3

G is constant along any leaf $\mathcal{S}_x$ of the foliation $\{\mathcal{S}_x\}$.

To proof this we define a basis of vector fields $(Y, f, \bar{f}, X, e, \bar{e})$, which is the respective dual of $(\Lambda, F, \bar{F}, T, E, \bar{E})$. Then we notice that the distribution $\mathcal{V}$ is spanned by vector fields of the form

$$V = Ue + \bar{U}\bar{e},$$

where U is any complex function on $\mathcal{U}$. Using an arbitrary V and the explicit form of G one easily finds that $\mathcal{L}_V G = 0$ due to the equations (1) and the properties of the

Lie derivative $\mathcal{L}_V$. $\square$

Now, we introduce an equivalence relation $\sim$ on $\mathcal{U}$ which identifies points lying on the same leave of $\{\mathcal{S}_x\}$. We assume that the quotient space $\mathcal{M} = \mathcal{U}/\sim$ is a 4-manifold. According to Lemma 3, G projects down to a well defined nondegenerate Lorentzian metric g on $\mathcal{M}$.

Theorem 1

The Lorentzian metric g on $\mathcal{M} = \mathcal{U}/\sim$ satisfies the Einstein equations $R_{ij} = \lambda g_{ij}$ and is not conformally flat.

Proof.

To proof the theorem, we first consider a 4-dimensional submanifold $\mathcal{M}'$ of $\mathcal{U}$ that is transversal to the leaves of $\{\mathcal{S}_x\}$. We have a natural inclusion $\iota : \mathcal{M}' \hookrightarrow \mathcal{U}$. $\mathcal{M}'$ may be equipped with a Lorentzian metric $g' = \iota^*G$. It is clear that $(\mathcal{M}, g)$ and $(\mathcal{M}', g')$ are locally isometrically equivalent. Thus, we may represent $(\mathcal{M}, g)$ by $(\mathcal{M}', g')$. In this way, it is enough to show that g' satisfies the Einstein equations. To do this, we first observe that $g' = \iota^*(F)\iota^*(\bar{F}) - \iota^*(T)\iota^*(\Lambda)$. Thus, a set of forms $\theta^i = (\iota^*(F), \iota^*(\bar{F}), \iota^*(T), \iota^*(\Lambda))$, ($i=1,2,3,4$) constitutes a null cotetrad for g' . To calculate the curvature we need to know $d\theta^i$. But these, due to the fact that $\iota^*d = d\iota^*$, are given by the realtions (1). To simplify the notation, we will omit the signs of the pullback ι^* in all of the formulae. On doing that we find that the connection 1-forms for g' determined by the equations

$$\begin{aligned} dF &= -\Gamma^1_1 \wedge F - \Gamma^1_3 \wedge T - \Gamma^1_4 \wedge \Lambda \\ d\bar{F} &= -\Gamma^2_2 \wedge \bar{F} - \Gamma^2_3 \wedge T - \Gamma^2_4 \wedge \Lambda \\ dT &= -\Gamma^3_1 \wedge F - \Gamma^3_2 \wedge \bar{F} - \Gamma^3_3 \wedge T \\ d\Lambda &= -\Gamma^4_1 \wedge F - \Gamma^4_2 \wedge \bar{F} - \Gamma^4_4 \wedge \Lambda \end{aligned}$$

$$g_{ik}\Gamma_j^k + g_{jk}\Gamma_k^i = 0, \quad g_{ij} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{pmatrix}$$

read

$$\Gamma^i_j = \begin{pmatrix} \bar{\Omega} - \Omega & 0 & -E & -\Gamma \\ 0 & \Omega - \bar{\Omega} & -\bar{E} & -\bar{\Gamma} \\ -\bar{\Gamma} & -\Gamma & \Omega + \bar{\Omega} & 0 \\ -\bar{E} & -E & 0 & -\Omega - \bar{\Omega} \end{pmatrix}.$$

The curvature $\mathcal{R}^i_j = d\Gamma^i_j + \Gamma^i_k \wedge \Gamma^k_j$ of this connection can be calculated using the relations (1), (2). Modulo the obvious reality conditions its components read as follows.

$$\begin{aligned}\mathcal{R}^1_3 &= \alpha F \wedge \Lambda - \bar{F} \wedge T \\ \mathcal{R}^1_4 &= -\bar{\alpha} T \wedge F + \bar{a}(\Lambda \wedge T + F \wedge \bar{F}) + h\bar{F} \wedge \Lambda \\ \frac{1}{2}(\mathcal{R}^3_3 + \mathcal{R}^1_1) &= (\bar{\alpha} + \frac{1}{2}\lambda)(\Lambda \wedge T + F \wedge \bar{F}) + \bar{a}\Lambda \wedge \bar{F}.\end{aligned}$$

These should be compared with the definitions of the spinorial coefficients of the Weyl tensor Ψ_μ , $\mu = 0, 1, 2, 3, 4$, the traceless Ricci tensor S_{ij} and the Ricci scalar R given by

$$\begin{aligned}\mathcal{R}^1_3 &= \bar{\Psi}_4 \bar{F} \wedge T + \bar{\Psi}_3(\Lambda \wedge T - F \wedge \bar{F}) + (\bar{\Psi}_2 + \frac{1}{12}R)\Lambda \wedge F \\ &\quad + \frac{1}{2}S_{33}F \wedge T + \frac{1}{2}S_{32}(\Lambda \wedge T + F \wedge \bar{F}) + \frac{1}{2}S_{22}\Lambda \wedge \bar{F},\end{aligned}$$

$$\begin{aligned}\mathcal{R}^1_4 &= (-\Psi_2 - \frac{1}{12}R)F \wedge T - \Psi_1(\Lambda \wedge T + F \wedge \bar{F}) - \Psi_0\Lambda \wedge \bar{F} \\ &\quad - \frac{1}{2}S_{22}\bar{F} \wedge T - \frac{1}{2}S_{42}(\Lambda \wedge T - F \wedge \bar{F}) - \frac{1}{2}S_{44}\Lambda \wedge F,\end{aligned}$$

$$\begin{aligned}\frac{1}{2}(\mathcal{R}^3_3 + \mathcal{R}^1_1) &= -\Psi_3 F \wedge T + (-\Psi_2 + \frac{1}{24}R)(\Lambda \wedge T + F \wedge \bar{F}) - \Psi_1 \Lambda \wedge \bar{F} \\ &\quad - \frac{1}{2}S_{32}\bar{F} \wedge T - \frac{1}{4}(S_{12} + S_{34})(\Lambda \wedge T - F \wedge \bar{F}) - \frac{1}{2}S_{41}\Lambda \wedge F.\end{aligned}$$

Looking at these equations we easily get $S_{ij} = R_{ij} - \frac{1}{4}Rg_{ij} \equiv 0$. This proves the Einstein property of the metric.

The spinorial coefficients for the Weyl tensor are also easy to obtain. They read

$$\Psi_0 = h, \quad \Psi_1 = -\bar{a}, \quad \Psi_2 = -\bar{\alpha} - \frac{\lambda}{3}, \quad \Psi_3 = 0, \quad \Psi_4 = -1.$$

Due to the nonvanishing of Ψ_4 the metric is never conformally flat.

We can also interpret the constant λ which for the first time appeared in equations (2). One easily reads from the above that it is proportional to the Ricci scalar $R = 4\lambda$. This concludes the proof of the Theorem. $\square$

We, additionally, note that the metric is of the Cartan-Petrov-Penrose type D iff

$$a = 0, \quad \text{and} \quad h = -(3\bar{\alpha} + \lambda)^2 \quad (8)$$

and of the type N iff

$$h = a = 0, \quad \text{and} \quad \lambda = -3\bar{\alpha}. \quad (9)$$

The metric is algebraically special iff $I^3 = 27J^2$, where

$$I = -h + \frac{1}{3}(3\bar{\alpha} + \lambda)^2, \quad \text{and} \quad J = \bar{a}^2 + \frac{1}{27}(3\bar{\alpha} + \lambda)^3 + \frac{1}{3}h(3\bar{\alpha} + \lambda).$$

The converse of the present paper can be stated in the following theorem (see Ref. [3] for details).

Theorem 2

Let $\mathcal{M}$ be a 4-dimensional manifold with a Lorentzian metric g satisfying the Einstein equations $R_{ij} = \lambda g_{ij}$. Then, there exists a double branched cover $\tilde{\mathcal{P}}$ of the bundle of null directions $\mathcal{P}$ over $\mathcal{M}$ [4, 5]

- i) which is a fibration over $\mathcal{M}$ with fibers being 2-dimensional tori (or, in algebraically special cases, their degenerate counterparts [3])*

and

- ii) on which there exists a unique system of 1-forms $(\Lambda, F, \bar{F}, T, E, \bar{E})$ which satisfies the following three conditions*

- i) Λ, T are real- and F, E are complex-valued 1-forms*
- ii) $\Lambda \wedge F \wedge \bar{F} \wedge T \wedge E \wedge \bar{E} \neq 0$ at each point p of $\tilde{\mathcal{P}}$*
- iii)*

$$\begin{aligned} dF &= (\Omega - \bar{\Omega}) \wedge F + E \wedge T + \Gamma \wedge \Lambda \\ dT &= \bar{\Gamma} \wedge F + \Gamma \wedge \bar{F} - (\Omega + \bar{\Omega}) \wedge T \\ d\Lambda &= \bar{E} \wedge F + E \wedge \bar{F} + (\Omega + \bar{\Omega}) \wedge \Lambda \\ dE &= 2\Omega \wedge E + \bar{F} \wedge T + \alpha \Lambda \wedge F. \end{aligned}$$

with certain complex-valued 1-forms Ω and Γ , and a certain complex function α on $\tilde{\mathcal{P}}$.

Acknowledgements

I warmly thank Andrzej Trautman for stimulating my interest in the papers [1, 2, 6]. I am also very grateful to Mike Crampin, Simonetta Frittelli, Jerzy Lewandowski, Lionel Mason, Ted Newman, David Robinson and Helmuth Urbantke for helpful discussions.

References

- [1] Cartan E 1932 “Sur la geometrie pseudo-conforme des hypersurfaces de deux variables complexes I” *Ann. Math. Pura Appl.* **11** 17-90
- [2] Cartan E 1932 “Sur la geometrie pseudo-conforme des hypersurfaces de deux variables complexes II” *Ann. Scuola Norm. Sup. Pisa* **1** 333-354
- [3] Nurowski P 1997 “Elliptic fibrations over the Einstein spacetimes” in preparation
- [4] Nurowski P 1996 “Optical geometries and related structures” *Journ. Geom. Phys.* **18** 335-348
- [5] Nurowski P 1997 “Twistor bundles, Einstein equations and real structures” in *Class. Q. Grav.* **14**, *The special volume edited by R. Penrose, I. Robinson and J. Tafel to celebrate A. Trautman’s 64-th birthday*, A261-A290.
- [6] Penrose R 1967 “The twistor programme” *Rep. Math. Phys.* **12** 65-76